

VU Research Portal

Opportunities and pitfalls in colorectal cancer screening

Terhaar sive Droste, J.S.

2011

document version

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

citation for published version (APA)

Terhaar sive Droste, J. S. (2011). *Opportunities and pitfalls in colorectal cancer screening*. [PhD-Thesis - Research and graduation internal, Vrije Universiteit Amsterdam].

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl



CHAPTER 6

Colonoscopy-controlled intra-individual comparisons to screen relevant neoplasia: Faecal Immunochemical Test versus Guaiac-based Faecal Occult Blood Test

J.S. Terhaar sive Droste [1], FA. Oort [1], R.W.M. van der Hulst [2], H.A. van Heukelem [3], R.J.L.F. Loffeld [4], I.C.E. Wesdorp [5], R.L.J. van Wanrooij [1], L. de Baaij [1], E.R. Mutsaers [1], S. van der Reijt [1], V.M.H. Coupe [6], J. Berkhof [6], A.A. Bouman [7], G.A. Meijer [8], C.J.J. Mulder [1]

[1] Gastroenterology and Hepatology, VU University Medical Center, Amsterdam

[2] Gastroenterology and Hepatology, Kennemer Gasthuis, Amsterdam

[3] Gastroenterology and Hepatology, Slotervaart Hospital, Amsterdam

[4] Internal Medicine, Zaans Medical Center, Zaandam

[5] Gastroenterology and Hepatology, Sint Lucas Andreas Hospital, Amsterdam

[6] Epidemiology and Biostatistics, VU University Medical Center, Amsterdam

[7] Clinical Chemistry, VU University Medical Center, Amsterdam

[8] Pathology, VU University Medical Center, Amsterdam

Abstract

Introduction: Guaiac-based Faecal Occult Blood Tests (g-FOBTs) are most commonly used in colorectal cancer (CRC) screening programs. Faecal Immunochemical Tests (FITs) are thought to be superior.

Methods: We assessed both sensitivity and specificity in different stages of CRC in eligible subjects who performed both FIT (OC-sensor®) and g-FOBT (Hemoccult-II®) prior to their scheduled colonoscopies.

Results: Of the 62 invasive cancers detected in 1821 individuals, g-FOBT was positive in 46 and FIT in 54 (74.2% versus 87.1%, $P=0.02$). Among 194 patients with advanced adenomas, g-FOBT was positive in 35 and FIT in 69 (18.0% versus 35.6%, $P<0.001$). 28 cancers were AJCC stage I or II, 31 stages III or IV, and in 3 cases stage was indeterminate. Sensitivity for screen relevant tumors (197 advanced adenomas and 28 stage I or II cancers) was 23.0% for g-FOBT and 40.5% for FIT ($P<0.001$). Specificity of g-FOBT compared to FIT for the detection of cancer was 95.7% versus 91.0% ($P<0.001$), and 97.4% versus 94.2% for advanced adenomas ($P<0.001$).

Discussion: FIT is more sensitive for CRC and advanced adenomas compared to g-FOBT. Particularly, sensitivity of FIT for screen relevant tumors, early stage cancers and advanced adenomas, is significantly higher. Specificity of g-FOBT is higher compared to FIT.

Introduction

Colorectal cancer (CRC) is a leading cause of cancer death world wide. Early detection is one of the most realistic approaches to reduce CRC-related death. Guaiac-based faecal occult blood tests (g-FOBTs) were already proposed for this purpose in the early 1970s.(1) Blood, shed into the colonic lumen by colorectal adenomas and carcinomas yields a positive g-FOBT due to the peroxidase-like activity of haeme in stool.(2) Screening programs using g-FOBT have proven to reduce both incidence and mortality of CRC.(3-7) Yet, both clinical sensitivity (i.e. the percentage of tumours detected in a series of tumor positive patients that perform the test) and program sensitivity (i.e. the percentage of tumours present in a population intended to screen that actually is detected) are suboptimal.(8,9)

More recently, the Faecal Immunochemical Test (or FIT), has been introduced as an alternative to g-FOBT. The FIT selectively detects the human globin-protein in stool, making it specific to colonic blood loss, while globin from blood lost proximal to the colon will be degraded before entering the colon. (10,11) Several variants of FIT exist, some of which come with automated analysis and have quantitative outcomes, like the one used in the present study.(12)

Comparisons of different techniques to detect occult blood in stool have been performed since 1953.(13) Recent studies that compared g-FOBT and FIT in screening populations, indicated superiority of FIT for the detection of both cancers and advanced adenomas.(8,14,15) In order to evaluate whether FIT can replace the most commonly used test (g-FOBT) in CRC screening, a comparative study design is needed. Both g-FOBT and FIT should be performed in parallel on the same stool samples.(14-16)

In addition, in order to appraise specificity of a test directly, all test negative individuals should undergo the test that is considered the gold standard, colonoscopy. Inherent to the design of screening studies, only FOBT-positive individuals underwent colonoscopy.(8,14,15,17) A large scale comparison of g-FOBT and FIT in a colonoscopy controlled population is still lacking. In the present study, test characteristics of both tests could be determined directly, as colonoscopy was performed in all included participants.

While population based screening studies yield crucial information on program sensitivity and acceptance of a test in the target population, often only small numbers of colorectal cancers are detected.(18) Consequently, the power to stratify these cancers by stage is insufficient. (8,19) In a referral population, like in the present study, a higher prevalence of CRC and its precursor lesions will allow for stratification of FOBT result for different phases of the natural history of the disease.

Aim

Aim of the present study was an intraindividual comparison of test performance of a g-FOBT and a quantitative FIT for detection of colorectal cancers of all stages and advanced adenomas in a colonoscopy controlled population. In addition, a specific aim was to compare test performance of both tests for the detection of early stage cancers and advanced adenomas taken together, since these lesions are most relevant for screening .

Methods

Study population

Five hospitals in the Amsterdam area in The Netherlands participated in this study. From June 2006 to March 2008, all ambulatory patients ≥18 years, scheduled for colonoscopy were invited to participate in this study, regardless of the indication for colonoscopy. Two of these five participating hospitals are situated in rural areas, another two are large teaching hospitals with an urban population. One of the centres is an academic medical centre with a predominantly urban population. In all centers, local Medical Ethics Review Board approval was obtained prior to the start of the study, and informed consent was obtained from the participants.

Inclusion and exclusion criteria

All eligible individuals were asked to perform both types of FOBT in the week preceding colonoscopy. Neither patients who presented at an emergency room nor institutionalized patients were enrolled. All indications for colonoscopy, as stated by the referring physicians, were recorded. Patients with a documented history of Inflammatory Bowel Disease (IBD) or with an incomplete colonoscopy were excluded from further analysis, as were patients who failed to complete both tests.

Study design

Elective patients were invited to participate either by their gastroenterologist when visiting the outpatient clinic, or by telephone by one of five researchers based at each of the participating centres. Once a patient consented in participation, an envelop was sent to his/her home address containing background information on the study, both FOBTs with extensive instructions and an informed consent form. When a person could not be reached by telephone, the same package was sent but with an additional explanatory letter.


FOBTs

The tests compared were the widely used guaiac based FOBT; Hemoccult II® (Beckman Coulter Inc., Fullerton, CA, USA) and an automated quantitative faecal immunochemical test: OC-sensor® (Eiken Chemical Co., Tokyo, Japan).

Faecal sampling

The three test-cards of the g-FOBT were sampled with stool from three separate bowel movements over a period of one week, 7, 3 and 1 day(s) prior to colonoscopy, respectively (figure 1). On the g-FOBT test cards, two separate samples from different parts of the stool had to be applied using the enclosed cardboard sticks. The final test card for g-FOBT and the FIT were sampled with stool taken from the day before colonoscopy, and before bowel preparation had started. Illustrated instructions guided the participants to sample their stool ensuring that contact with water or urine was prevented. No restrictions were made with regard to either diet or use of medication during the week in which stool samples were taken.(20)

Figure 1: Study design



Test analysis

On the day of colonoscopy, both the completed FOBTs and the informed consent form were handed over to the nursing staff at the endoscopy-department. All FITs were stored at minus 5 degrees Celsius on arrival and all g-FOBTs were stored at room temperature according to

the manufacturers recommendations. Tests were analyzed according to the manufacturers instructions within one week by two experienced technicians, who were unaware of the clinical data.(21) Both technicians were tested negative for colour blindness and received special training for reading the test cards.

FIT samples were processed with the OC sensor MICRO desktop analyser (Eiken Chemical co, Tokyo, Japan).(21) A haemoglobin concentration of $\geq 100\text{ng/ml}$ was taken as cut off, according to the manufacturers recommendations. G-FOBT samples were considered positive when a blue colour appeared in at least 1 out of 3 test cards, following application of the reagent. Test cards were not rehydrated prior to analysis.

Colonoscopy

All participants underwent complete colonoscopy in one of five participating centres. Colonoscopies were performed, or supervised, by experienced gastroenterologists. Conscious sedation using Midazolam was offered to all patients. Endoscopists were blinded to the results of both tests. A complete colonoscopy was defined as intubation of the caecum with identification of the ileocecal valve or appendiceal orifice, or intubation up to an obstructing neoplasm. In addition, patients with inadequate bowel cleansing, as judged by the individual endoscopist, were excluded as well. Patients were classified based on the most advanced lesion detected in their colon. Estimated size of all lesions, as reported in the colonoscopy report, were categorized.

Detected lesions

Histology of tissue samples obtained during colonoscopy was evaluated according to standard procedures. Pathology reports were collected centrally and entered into the database. Adenomas $\geq 1.0\text{ cm}$, with any villous features (i.e. tubulovillous or villous adenoma) or high-grade dysplasia, were considered advanced adenomas.(22,23) Advanced neoplasia included all cases of colorectal cancer and all advanced adenomas. Colorectal carcinomas were staged according to the AJCC cancer staging manual.(24)

Statistical analysis

Defining colonoscopy outcome as gold standard, sensitivity and specificity of both FOBTs were calculated, for advanced adenomas and for CRC separately. Furthermore, these test characteristics were calculated for all participants with either one or more advanced adenoma(s) or an early stage carcinoma (AJCC stage I and II). Since both FOBTs were performed in parallel on the same stool samples, McNemar’s test could be used for comparison of correlated proportions.

The sample size was predetermined, based on an anticipated difference in sensitivity for detection of colorectal cancer of 20% between Hemoccult II and OC-sensor. The enrolment

of 1650 participants was required, assuming a prevalence of colorectal cancer of at least 2,5% in this study population, to provide the study with a statistical power of 80% to detect a significant difference at a two-sided alpha level of 0.05 with the use of McNemar’s test. All analyses were performed with SPSS for Windows Version 15 (SPSS Inc., Chicago, USA).

Results

Demographics

Overall 2217 individuals who underwent colonoscopy performed at least one of both FOBTs. In total 396 individuals were excluded for reasons listed in Table 1. The mean age of 1821 individuals that were included for final analysis was 59.6 years (range 18-86 years, 95% CI 59.0- 60.1). The majority of the subjects (56.9%) was female.

Indications for colonoscopy

The primary indication for colonoscopy was available for 1742 out of 1821 individuals. In 79 patients (4.3%), the primary indication remained unspecified. Indications were classified in four categories listed in Table 2. Almost half of the population (N=897, 49.3%) was referred for colonoscopy because of gastrointestinal symptoms and 44.3% of patients was asymptomatic and had an indication for screening or surveillance colonoscopy (N=807).

Table 1: Reasons for exclusion of 396 patients out of 2221 consecutive patients undergoing colonoscopy in five hospitals in the Amsterdam area.

Reason for exclusion		N=396 (100%)
Incomplete colonoscopy	Caecum not reached	102 (25.8%)
	Insufficient bowel lavage	27 (6.8%)
Documented history of IBD		77 (19.4%)
FIT not sampled		87 (22.0%)
g-FOBT not sampled		98 (24.7%)
Colostomy		5 (1.3%)

Table 2: Primary indications for colonoscopy among 1821 consecutive patients in five hospitals in the Amsterdam area enrolled in a study comparing FIT versus g-FOBT

Indication Group	Indication for colonoscopy	N
Symptomatic/suspect		
	Weightloss	15
	Clinical suspicion of diverticulitis	14
	Clinical suspicion of IBD	17
	Abdominal pain	280
	Anemia	77
	Hematochezia	190
	Altered bowel habits	304
	Clinical suspicion of CRC (inconclusive pathology)	1
	Colonoscopy for polypectomy	37
	Total	935
Screening & Surveillance		
	Average risk	32
	Familial history of CRC	288
	Lynch syndrome	29
	Polyp surveillance	314
	Post CRC surveillance	120
	Radiological suspicion of malignancy	21
	Total	804
Other		
	Not specified	82
	Grand total	1821

Colonoscopy results

In 194 of 1821 patients (10.7%), at least one advanced adenoma was found on colonoscopy. Adenocarcinomas were found in 62 of 1821 patients (3.4%). Of these, 28 (45.2%) were classified as early stage colorectal cancer (AJCC stage I and II) and 31 patients (50.0%) as late stage (AJCC stage III and IV). Three rectal cancers could not be accurately staged due to the effects of neo-adjuvant radiotherapy.

Test results

Overall positivity rate for the g-FOBT was 6.7% (122/1821) and for the FIT 11.8% (214/1821). The g-FOBT detected 46 of 62 cancers for a sensitivity of 74.2%, whereas the FIT detected 54 of 62

cancers, for a sensitivity of 87.1% (Table 3). FIT detected 7 cancers that did not score a positive g-FOBT. Only one cancer scored a positive g-FOBT but not a positive FIT. The observed difference in sensitivities between the two tests was significant ($P=0.02$). Sensitivity for screen relevant lesions (advanced adenomas and early stage cancers) was 23.0% for g-FOBT and 40.5% for FIT ($P<0.001$).⁽²⁵⁾ Yet, g-FOBT was found to have a higher specificity compared to FIT for cancers of all stages (95.7% vs. 91.0%, $P<0.001$) and for advanced adenomas (97.4% vs. 94.2%, $P<0.001$). The difference between sensitivities of g-FOBT and FIT for the detection of early or late stage colorectal cancers respectively, showed a similar trend but did not reach statistical significance. The sensitivities of g-FOBT and FIT for early stage cancers was 57.1% and FIT 75.0%, respectively (Table 4). For late stage cancers g-FOBT had a sensitivity of 87.1% and FIT 96.8%.

Table 3: Sensitivity and specificity of g-FOBT (Hemoccult II®) and FIT (OC-sensor®)[§] for detection of advanced adenomas, advanced neoplasia, colorectal cancer (CRC), in a consecutive series of 1821 patients referred for colonoscopy.

	Advanced Adenomas	Early stage CRC (Stage I&II) + Advanced Adenomas*	CRC	Advanced Neoplasia
	194/1821	222/1821	62/1821	256/1821
Sensitivity Hemoccult II® (95% CI)	18.0 % 35/194 (12.9–24.2)	23.0 % 51/222 (17.6–29.1)	74.2 % 46/62 (61.5–84.5)	31.6 % 81/256 (26.0–37.7)
Sensitivity OC-sensor® (95% CI)	35.6 % 69/194 (28.9–42.7)	40.5 % 90/222 (34.1 – 47.3)	87.1 % 54/62 (76.2–94.3)	48.1 % 123/256 (41.8–54.4)
p-value**	<0.001	<0.001	0.02	<0.001
Specificity Hemoccult II® (95% CI)	97.4 % 1524/1565 (96.5–98.1)	97.4 % 1524/1565 (96.5–98.1)	95.7 % 1683/1759 (94.6–96.6)	97.4 % 1524/1565 (96.5–98.1)
Specificity OC-sensor® (95% CI)	94.2 % 1474/1565 (92.9–95.3)	94.2 % 1474/1565 (92.9–95.3)	91.0 % 1599/1759 (89.5–92.2)	94.2 % 1474/1565 (92.9–95.3)
p-value**	<0.001	<0.001	<0.001	<0.001

*Of three rectal tumours the oncological stage of disease could not be assessed due to the effects of neo-adjuvant radiotherapy.

**McNemar's test was used to compare paired proportions

[§] cut-off level ≥ 100 ng/ml

Discussion

In the present study, performance characteristics of both g-FOBT and FIT were evaluated in parallel in a referral population of 1821 patients who underwent complete colonoscopy. This design allowed to deal with two major issues that, thus far, have remained unaddressed. Firstly, in most studies so far on FIT, only individuals with a positive test underwent subsequent colonoscopy, which precluded the determination of false negativity rates and thus specificity of the investigated tests. Secondly, previous studies comparing g-FOBT and FIT did so mainly in separate patient groups, which limits determination of exactly which tumors are missed by one test but are detected by the other.

Overall, results of the present study are consistent with earlier observations in screening studies and confirm a significantly higher sensitivity of FIT for colorectal cancer as well as advanced adenomas. The referral population that was evaluated in the present study consisted of more individuals with CRC or advanced adenomas than an average risk screening population. Due to the relatively high tumor yield in the referral population in the present study, test results of FIT and g-FOBT could be stratified for early stage cancers (AJCC Stage I and II) and late stage cancers (Stage III and IV). Importantly, for the screen relevant neoplasia (i.e. advanced adenomas and early stage cancers taken together), FIT significantly outperforms g-FOBT in terms of sensitivity with 40.5% versus 23.0%, respectively. This is of special importance since population-based screening programs for CRC aim to detect this category of neoplasia

Table 4: Test outcome of g-FOBT (Hemoccult II®) and FIT (OC-sensor®)§ in 62 patients diagnosed with colorectal cancer at an early stage (n=28) and at a late stage (n=31)

	Hemoccult II®		OC-sensor®		Total
	Positive	Negative	Positive	Negative	
Early stage disease (Stage I & II)	16 (57.1%)	12 (42.9%)	21 (75.0%)	7 (25.0%)	28
Late stage disease (Stage III and IV)	27 (87.1%)	4 (12.9%)	30 (96.8%)	1 (3.0%)	31
AJCC stage unknown*	3	0	3	0	3
Total	46 (74.2%)	16 (25.8%)	54 (87.1%)	8 (12.9%)	62 (100%)

* three rectal tumours could not be accurately staged due to the effects of neo-adjuvant radiotherapy.
§ cut-off level ≥ 100 ng/ml

specifically. A trend in the same direction was observed in early stage cancers alone, with detection rates of 75.0% versus 57.1% for FIT and g-FOBT, respectively, but this did not reach statistical significance.

In a screening-naïve population both early and late stage colorectal cancers, the so called prevalent cancers, will be detected in a range consistent with their respective prevalences. Therefore, in a first round of a screening program, overall CRC detection rates will be inflated by the prevalent advanced stage cancers. However, given the nature of the disease, with usual annual or biennial screening programs, in a second round of screening, less advanced cancers will be left in the population.(26) Therefore, the performance characteristics of the screening program will largely depend on the potential to detect early stage CRCs, i.e. incident cancers. In this respect, it is highly relevant that the present study allowed to analyze performance of both tests separately for early and late stage CRC. Preferably, these results need to be validated in a screening population, but the small numbers of CRCs detected in screening studies will hamper such a study design.

Positive and negative predictive values of a test are influenced by the prevalence of a disorder in the population that is tested. Therefore, in this referral population, these values were not calculated as they would not reflect predictive values in a screening population. However, sensitivity and specificity are characteristics of a diagnostic test and are not influenced by the prevalence of a disease in population.(27)

With respect to whether the present findings are generalizable to a screening population, it could be argued that preclinical (i.e. screen detected lesions) may be different (e.g. in their tendency to bleed) from symptomatic lesions. However, so far there is no evidence that either supports or falsifies this hypothesis.

The present study shows that the higher sensitivity of FIT goes at the cost of a somewhat lower specificity (advanced neoplasia; 97.4% versus 94.2%, overall cancer; 95.7% versus 91.0%, for g-FOBT and FIT, respectively). The higher specificity of g-FOBT compared to FIT, was recently described in another study comparing exactly the same g-FOBT and FIT, but in two separate populations.(8) In this study, that used a population screening design, specificity could not be calculated directly, since only FIT or g-FOBT positive individuals were offered colonoscopy. Hence specificities were calculated based on rare disease assumptions. This may explain the small differences in specificity between their study and the current one, in which all individuals underwent colonoscopy. Moreover, not all guaiac based FOBTs have the same test characteristics. The Hemoccult II used in the present study is a low sensitivity FOBT. The United States only endorse the use of sensitive g-FOBTs (e.g. Hemoccult Sensa) for screening and therefore the results might be different if an FOBT with better sensitivity was used.(28)

Apart from the higher sensitivity for screen relevant tumors, FIT has several other advantages. The technical characteristics of the FIT test used in the present study allow for automated analysis, unlike g-FOBT, which makes it suitable for high throughput application. Furthermore, patient acceptance of a test is a major determinant of compliance and consequently success of a screening program. The OC-sensor® has been found to be significantly better accepted by the average risk target population for colorectal cancer screening than the Hemoccult II®.(8) Finally, the reproducibility and quality control of FIT is good, where g-FOBT was not optimal.(29,30)

Secondary prevention of CRC is a major health care issue, and several countries already have introduced g-FOBT in large pilot-studies or nationwide colorectal cancer screening programmes.(31,32) A growing body of literature lends support to the notion that FIT is superior to g-FOBT in colorectal cancer screening.(8,15,17,33,34) The present study adds to this the observation that FIT has a significantly higher sensitivity for screen relevant tumors than g-FOBT. In addition, the present study allows for a more precise estimation of the specificity of FIT for colorectal tumors.

References

1. Gregor DH. A progress report. Detection of colorectal cancer using guaiac slides. *CA Cancer J.Clin.* 1972; 22: 360-363. Ref Type: Journal
2. Allison JE. Colon Cancer Screening Guidelines 2005: the fecal occult blood test option has become a better FIT. *Gastroenterology* 2005; 129: 745-748. Ref Type: Journal
3. Mandel JS, Church TR, Bond JH, Ederer F, Geisser MS, Mongin SJ, Snover DC, Schuman LM. The effect of fecal occult-blood screening on the incidence of colorectal cancer. *N.Engl.J.Med.* 2000; 343: 1603-1607. Ref Type: Journal
4. Kronborg O, Fenger C, Olsen J, Jorgensen OD, Sondergaard O. Randomised study of screening for colorectal cancer with faecal-occult-blood test. *Lancet* 1996; 348: 1467-1471. Ref Type: Journal
5. Mandel JS, Bond JH, Church TR, Snover DC, Bradley GM, Schuman LM, Ederer F. Reducing mortality from colorectal cancer by screening for fecal occult blood. Minnesota Colon Cancer Control Study. *N.Engl.J.Med.* 1993; 328: 1365-1371. Ref Type: Journal
6. Hardcastle JD, Chamberlain JO, Robinson MH, Moss SM, Amar SS, Balfour TW, James PD, Mangham CM. Randomised controlled trial of faecal-occult-blood screening for colorectal cancer. *Lancet* 1996; 348: 1472-1477. Ref Type: Journal
7. Allison JE. Review article: faecal occult blood testing for colorectal cancer. *Aliment.Pharmacol.Ther.* 1998; 12: 1-10. Ref Type: Journal
8. van Rossum LG, van Rijn AF, Laheij RJ, van Oijen MG, Fockens P, van Krieken HH, Verbeek AL, Jansen JB, Dekker E. Random Comparison of Guaiac and Immunochemical Fecal Occult Blood Tests for Colorectal Cancer in a Screening Population. *Gastroenterology* 2008; 135: 82-90. Ref Type: Journal
9. Young GP, Sinatra MA, St John DJ. Influence of delay in stool sampling on fecal occult blood test sensitivity. *Clin.Chem.* 1996; 42: 1107-1108. Ref Type: Journal
10. Young GP, Cole S. New stool screening tests for colorectal cancer. *Digestion* 2007; 76: 26-33. Ref Type: Journal
11. Kronborg O, Regula J. Population screening for colorectal cancer: advantages and drawbacks. *Dig.Dis.* 2007; 25: 270-273. Ref Type: Journal
12. Launoy GD, Bertrand HJ, Berchi C, Talbourdet VY, Guizard AV, Bouvier VM, Caces ER. Evaluation of an immunochemical fecal occult blood test with automated reading in screening for colorectal cancer in a general average-risk population. *Int.J.Cancer* 2005; 115: 493-496. Ref Type: Journal
13. HEPLER OE, WONG P, PIHL HD. Comparison of tests for occult blood in feces. *Am.J.Clin.Pathol.* 1953; 23: 1263-1272. Ref Type: Journal

14. Allison JE, Sakoda LC, Levin TR, Tucker JP, Tekawa IS, Cuff T, Pauly MP, Shlager L, Palitz AM, Zhao WK, Schwartz JS, Ransohoff DF, Selby JV. Screening for colorectal neoplasms with new fecal occult blood tests: update on performance characteristics. *J.Natl.Cancer Inst.* 2007; 99: 1462-1470. Ref Type: Journal
15. Dancourt V, Lejeune C, Lepage C, Gailliard MC, Meny B, Faivre J. Immunochemical faecal occult blood tests are superior to guaiac-based tests for the detection of colorectal neoplasms. *Eur.J.Cancer* 2008; 44: 2254-2258. Ref Type: Journal
16. Bossuyt PM, Irwig L, Craig J, Glasziou P. Comparative accuracy: assessing new tests against existing diagnostic pathways. *BMJ* 2006; 332: 1089-1092. Ref Type: Journal
17. Guittet L, Bouvier V, Mariotte N, Vallee JP, Arsene D, Boutreux S, Tichet J, Launoy G. Comparison of a guaiac based and an immunochemical faecal occult blood test in screening for colorectal cancer in a general average risk population. *Gut* 2007; 56: 210-214. Ref Type: Journal
18. Ransohoff DF. What is the role of iFOBT in screening for colorectal cancer? *Gut* 2007; 56: 1343-1344. Ref Type: Journal
19. Young GP, Cole SR. Which fecal occult blood test is best to screen for colorectal cancer? *Nat. Clin.Pract.Gastroenterol.Hepatol.* 2009. Ref Type: Journal
20. Pignone M, Campbell MK, Carr C, Phillips C. Meta-analysis of dietary restriction during fecal occult blood testing. *Eff.Clin.Pract.* 2001; 4: 150-156. Ref Type: Journal
21. Rozen P, Waked A, Vilkin A, Levi Z, Niv Y. Evaluation of a desk top instrument for the automated development and immunochemical quantification of fecal occult blood. *Med. Sci. Monit.* 2006; 12: MT27-MT32. Ref Type: Journal
22. Rex DK, Kahi CJ, Levin B, Smith RA, Bond JH, Brooks D, Burt RW, Byers T, Fletcher RH, Hyman N, Johnson D, Kirk L, Lieberman DA, Levin TR, O'Brien MJ, Simmang C, Thorson AG, Winawer SJ. Guidelines for colonoscopy surveillance after cancer resection: a consensus update by the American Cancer Society and US Multi-Society Task Force on Colorectal Cancer. *CA Cancer J.Clin.* 2006; 56: 160-167. Ref Type: Journal
23. Winawer SJ, Zauber AG, O'Brien MJ, Ho MN, Gottlieb L, Sternberg SS, Wayne JD, Bond J, Schapiro M, Stewart ET, . Randomized comparison of surveillance intervals after colonoscopic removal of newly diagnosed adenomatous polyps. The National Polyp Study Workgroup. *N.Engl.J.Med.* 1993; 328: 901-906. Ref Type: Journal
24. Greene Fea. AJCC (American Joint Committee on Cancer) Cancer Staging Manual. Springer-Verlag, 2002: 113. Ref Type: Book Chapter
25. Ahlquist DA, Sargent DJ, Loprinzi CL, Levin TR, Rex DK, Ahnen DJ, Knigge K, Lance MP, Burgart LJ, Hamilton SR, Allison JE, Lawson MJ, Devens ME, Harrington JJ, Hillman SL. Stool DNA and occult blood testing for screen detection of colorectal neoplasia. *Ann.Intern.Med.* 2008; 149: 441-50, W81. Ref Type: Journal
26. Grazzini G, Ciatto S, Cislighi C, Castiglione G, Falcone M, Mantellini P, Zappa M. Cost evaluation in a colorectal cancer screening programme by faecal occult blood test in the District of Florence. *J.Med.Screen.* 2008; 15: 175-181. Ref Type: Journal
27. Altman D. Practical statistics for medical research. 409-413. 8 A.D. Chapman&Hall/CRC. Ref Type: Generic
28. Screening for colorectal cancer: U.S. Preventive Services Task Force recommendation statement. *Ann.Intern.Med.* 2008; 149: 627-637. Ref Type: Journal
29. Levi Z, Rozen P, Hazazi R, Vilkin A, Waked A, Maoz E, Birkenfeld S, Leshno M, Niv Y. A quantitative immunochemical fecal occult blood test for colorectal neoplasia. *Ann.Intern.Med.* 2007; 146: 244-255. Ref Type: Journal
30. Selinger RR, Norman S, Dominitz JA. Failure of health care professionals to interpret fecal occult blood tests accurately. *Am.J.Med.* 2003; 114: 64-67. Ref Type: Journal
31. Denis BJ, Ruetsch M, Strentz P, Vogel JY, Guth F, Boyaval JM, Pagnon X, Ebelin JF, Gendre I, Perrin P. Short-term outcomes of the first round of a pilot colorectal cancer screening programme with guaiac based faecal occult blood test. *Gut* 2007. Ref Type: Journal
32. Benson VS, Patnick J, Davies AK, Nadel MR, Smith RA, Atkin WS. Colorectal cancer screening: a comparison of 35 initiatives in 17 countries. *Int.J.Cancer* 2008; 122: 1357-1367. Ref Type: Journal
33. Wong BC, Wong WM, Cheung KL, Tong TS, Rozen P, Young GP, Chu KW, Ho J, Law WL, Tung HM, Lai KC, Hu WH, Chan CK, Lam SK. A sensitive guaiac faecal occult blood test is less useful than an immunochemical test for colorectal cancer screening in a Chinese population. *Aliment.Pharmacol.Ther.* 2003; 18: 941-946. Ref Type: Journal
34. Levi Z, Hazazi R, Rozen P, Vilkin A, Waked A, Niv Y. A quantitative immunochemical faecal occult blood test is more efficient for detecting significant colorectal neoplasia than a sensitive guaiac test. *Aliment.Pharmacol.Ther.* 2006; 23: 1359-1364. Ref Type: Journal